COMMUNICABLE DISEASE CENTER

POLIOMYELITIS

SURVEILLANCE

TABLE OF CONTENTS

SUMMARY

- I. CURRENT MORBIDITY TRENDS
- II. 1962 CASES REPORTED TO PSU
- III. ROUTINE SURVEILLANCE 1962 CASES
- IV. ENTEROVIRUS SURVEILLANCE
- V. FOREIGN REPORTS BRITISH GUIANA
- VI. STATEMENT ON THE USE OF OPV BY THE AMERICAN ACADEMY OF PEDIATRICS

SUPPLEMENT TO PSU 274 – Index to 1962 PSU Reports



National
Communicable Disease Center
Library
Atlanta, Georgia 30333

PREFACE

Summarized in this report is information received from State Health Departments, university investigators, virology laboratories and other pertinent sources, domestic and foreign. Much of the information is preliminary. It is intended primarily for the use of those with responsibility for disease control activities. Anyone desiring to quote this report should contact the original investigator for confirmation and interpretation.

Contributions to the Surveillance Report are most welcome. Please address to: Chief, Poliomyelitis Surveillance Unit, Communicable Disease Center, Atlanta 22, Georgia.

Communicable Disease Center

Epidemiology Branch

Statistics Section
Surveillance Section

Poliomyelitis Surveillance Unit

James L. Goddard, M.D., Chief

Alexander D. Langmuir, M.D., Chief

Robert E. Serfling, Ph.D., Chief Donald A. Henderson, M.D., Chief

James A. Bryan, II, M.D., Chief John J. Witte, M.D. Saul Rosenberg, Statistician

SUMMARY

The expected seasonal decline in cases has occurred with only 32 paralytic cases being reported during the four-week period ending January 12. These largely represent delayed reports with no case concentrations being noted.

The preliminary figures for paralytic and total cases of poliomyelitis for 1962 show a 20 percent decline in paralytic cases in comparison with the figures for 1961, the preceding record low year.

Individual case forms have been received on 653 of the 708 paralytic cases reported in 1962. Of these cases, 23 percent gave a history of having received 3 or more doses of inactivated vaccine. The reports of virologic studies received thus far on these cases are also shown in Section 2.

Surveillance data on cases occurring within 30 days of receiving the poliomyelitis vaccines are presented in Section 3. Sixteen paralytic cases have been noted following inactivated vaccine and 58 following oral vaccine.

Enterovirus isolates have been reported from 34 States with Coxsackie B-3 and ECHO 9 being reported from the greatest number of these States.

In Section 5 a preliminary report on the current poliomyelitis outbreak in British Guiana is presented. Three hundred sixty-one cases have been reported from November 1, 1962 through January 15, 1963.

The position taken by the American Academy of Pediatrics on the use of oral poliomyelitis vaccine is presented in Section 6.

The Index for 1962 Poliomyelitis Surveillance Unit Reports is appended as a supplement to this issue.

1. CURRENT POLIOMYELITIS MORBIDITY TRENDS

During the four weeks from December 16, 1962 through January 12, 1963, a total of 37 cases of poliomyelitis, 32 paralytic, were reported. Eighteen of the paralytic cases were reported for the week ending December 22.

Reports of cases during the past four weeks came from 14 States and were in the main delayed reports. Pennsylvania accounted for 7 paralytic cases while California and Louisiana reported 6 cases each.

The totals for cases reported by telegram through the last week in 1962 are shown below compared with those of the preceding four years.

Poliomyelitis (1st through 52nd Week) for the Past Five Years

	1962	<u>1961</u>	<u>1960</u>	1959	1958	
Paralytic	708	884	2292	5715	3150	
Total	889	1364	3301	8573	6092	

THAMMUE

2. 1962 POLIOMYELITIS CASES REPORTED TO PSU

A total of 889 cases of poliomyelitis with onset in 1962 has been reported by weekly telegrams to the CDC Morbidity and Mortality Unit. Of these, 708 have been paralytic. These figures represent a preliminary count. The final data will include 1962 delayed case reports and case revisions, and will be presented in the annual supplement of the Morbidity and Mortality Weekly Report in the latter part of this year.

The Poliomyelitis Surveillance Unit has received individual case forms on 653 of the 708 paralytic cases reported. The vaccination status of these 653 paralytic cases by age groups is shown below.

for 1962 show a 20 percent decline in paralytic cases in comparison with

adjusted to whose in a wag inserted () seems seem 10 .2001 at beinger seems algologic to at Paralytic Poliomyelitis by Age Grouped side to be viscer . I no And Vaccination History Reported on PSU Forms bevieves galbute (Through January 12, 1963)

lance data on cases occurring within 10 days of receiving the	Surveill
Ageness size a Doses of Inactivated Vaccine sassers are serious	
Group and I OV an 1-2V at 3V an 4+V at a Unk. several Total revolted I	Percent and avail
0-4022000 212ty se 5018 A16poxî 23: roge11cosd byed 312: sleak auxi	47.8
5-9 , ass 64 see 17to /122 num 32 see 3 3 mort 5138 que a guied	
10-14 23 15 18 9 3 68	
ion 5 a pre82minary replac on 4 he curbent p41iomye21tis ov91-24k	154.3 T
20-29 and 1381 ages 6 and 3 tolk b 4 about 2 will bes 53 earq at any	
30-39 21 6.882 through January 15, 2963. 6	1 m4.9 beiroger
40+ <u>18 1 0 2 1 22</u>	3.4
ltion taken by the American Academy of Pediatrics on the	
TOTAL 389 .99 m 1 673 mi 176 mes 22 el en lo 653 il	100.0 to each
ex for 1962 Policmyelitis Jurveillance Unit Reports is TMSDREY	
DOSES 61.7 15.7 10.6 12.0 - see a sed 100.0 seme focus	larkmanne

Of the 301 cases with known vaccination history in the 0-4 age group, 212 (70.4 percent) had not received any poliomyelitis vaccinations and only 39 (13.0 percent) gave a history of three or more doses of inactivated vaccine.

As of January 12, 1963, the Poliomyelitis Surveillance Unit has received results of virological studies on 406 of the 653 paralytic cases. A total of 350 poliovirus isolates (86.2 percent) has been recovered from the 406 cases. Of these, 265 are Type I, 5 are Type II, and 80 are Type III polioviruses. Isolates associated with clinical paralytic poliomyelitis have been reported from the following States.

Poliomyelliis (lat through 52md sex) for the east Five Years

	Po	liovirus		
State	<u> 1 962</u>	all call	Total	
Alabama	snippsv 1 11	0 0	Joans Cliv ass. 11	
Arizona	2	0 0	2	
Arkansas	.at we mend	1 0	8 Buring the four-or	
California	25	0 12	vilog la racso per 37 .	
Colorado	noon need 1	2 2	v salvolla) days days v Colboras (brass of the	13 1
	Surveillance	2.13 (1.59)10 (1.60	sentili to tosno bai	
D.C. Florida	acons, Califo	TA 0, 298 (18)	assumed to strain but	
correlation poits lerron	OH RAY BIY	0.55 5.25	nno red estavopos s2	
Idaho	21 Talk in the	0 1	oria has notroetal 50 s	
	20	-	for To maintail eath 30	
1 (///) 1		is vince value	in to hir total portr 20	
Previous Para,	abo Guset	0 0	esă vintet 6	
	13	0 0 1	14	
Kentucky Louisiana	1/	-	n especie 22	
	Ofto a care	0 0	s alter 1	
Maryland	27/23	And the second of the second		
Massachusetts	arab SI 3	0 7/6 ON	and Canthernon 18,122	
Michigan	1.80 vaccine c	0 4	r boshow a cafe sould be	
Minnesota	4	0 0	4	
Mississippi	i mer amar 1	2 to 0 a vino 17 cm	in sympo and Jawe 8	
	5	2 0	i už doblos sniemo os 4 .	
Montana	electros a d	0 0	-boyalibarre? Midutelmo	
Nebraska	O Paralyti	0 1	agrificationals broundles 1	
New Hampshire	Anney 90 Lu	0 2	John Charman .83	
New Jersey	The second secon	0 0	1	
Norr Morrison	rhanen en 2	0 0	rek salad Joyana 169 edil 2	
New York	, 200 f at -7	0 7	7, 10 : 10 10 20 10 11 14 10	
North Carolina	eu d'odo il e. La di dona 3 u	0) 100	tala ilmi yanan aka min Tamakiliko ni bamman 3 k	
Ohio	9	0 4	y action during a population	
Oregon	0	0 3	trug angab 0% pentitiva a3 in-	
Pennsylvania	5	0 5	10	
South Carolina	გესადეგ დად ა	0, 3	cell survey floret 5	
Tennessee	j-, a 3	91 0 0 an 1	4	
Texas	90	0 7	97 x 2 x 2 x 2 x 3 x 3 x 3 x 3 x 3 x 3 x 3	
Utah	1	0 0	1	
Washington	3	0 1	.e % 26/20 Lité 4	
West Virginia	10	0 1	O1 3 25 Fig. 11 s	
		1 0/23 1	of R Radio	
Wisconsin	gy03 € 1	1 (0)(0) 1 3	(1) No contract 2 >	
Wyoming	erisb 8 1	0 0	(0 f % 10 k <u>rs/1</u>	
momar a	265	5 80	350	
TOTALS	265	3 80	330	

3. ROUTINE POLIOMYELITIS SURVEILLANCE - 1962

A. Cases with Onset within 30 Days of Vaccination (Inactivated)

Illinois

During the four-week period from December 15, 1962 to January 12, 1963, three cases of poliomyelitis (including 2 paralytic) occurring within 30 days following vaccination have been reported on individual surveillance forms to the Poliomyelitis Surveillance Unit. All three cases had onset of illness in 1962. Alabama, California, and South Carolina accounted for one case each. There was no correlation between site of injection and site of first paralysis in the two paralytic cases.

A line listing of the three cases is given below:

State	d Ar	County	Age	Sex Onset	Date Vacc.	Onset Interval	Previous IPV	Para. Status Isol.
Alabama	53	Choctaw	6	M 9/17	9/15	2 days	•	sospaluod I
California;	+	Marin	33	F 10/22	10/17	5 days		NP
So. Carolin	ıa	Charleston	18/12	M 9/26	9/14	12 days	22339	I Massa q inus

^{*} This case also received Type I oral polio vaccine on 9-23.

Twenty-two cases of poliomyelitis (16 paralytic) with onset in 1962, occurring within 30 days of vaccination, have been reported to the Poliomyelitis Surveillance Unit. There was a correlation between site of injection and site of first paralysis in one paralytic case.

B. Cases with Onset within 30 Days of Vaccine (Oral) - 1962

The Poliomyelitis Surveillance Unit has received reports of 6 additional cases of poliomyelitis with onset in 1962, occurring within 30 days following oral vaccination. Five of the 6 were paralytic. Three cases occurred in California, and one case each in Alabama, Montana, and Texas. Type I oral polio vaccine was administered to each of the 6 patients within 30 days prior to onset.

A detailed line listing of the 6 cases appears below.

State	County	Age	Sex	Onset	Date Fed	Type Onset Fed Interval	Doses IPV	Para. Status Isolate
Alabama	Mobile 2	3/12	F	9/17	9/6	I 11 days	3	Mashing q an
California		3 3	F	10/22	9/23	I 29 days	5	NP
California	Marin	30	M	10/7	9/23	I 14 days	1	P III
California		35	M	11/6	10/28	I 9 days	0	th succes I
Montana	Park	39	F	10/16	10/7	I 9 days	3	B gurno Te
Texas	Ellis	1	\mathbf{F}	8/12	8/10	I 2 days	0	PI

Sixty-seven cases of poliomyelitis (58 paralytic) with onset in 1962, occurring within 30 days following oral vaccination have been submitted to the Poliomyelitis Surveillance Unit on individual surveillance forms. Of the paralytic cases, 31 had received Type I oral polio vaccine within 30 days prior to onset, 3 had taken Type II vaccine, and 24 had been fed Type III vaccine. From month of vaccine licensures through November 1962, 33 million doses of Type I oral polio vaccine, 14 million doses of Type II vaccine, and 12 million doses of Type III vaccine have been administered in known mass community oral vaccination programs.

In. 8.8.6. Nehaul, Acting Chick Medical Officer for Trivish Criang has recorded the congresse of 30 cases of paralytic pelcomposites through James, 5, recurring in an entirement which began during the very years lames, 100 percent of the age group involved has been predominantly the very years which SO percent of the patients being below six years of age and almost of percent being below 2 years of abo. Tyre I policyled has been incomence from a tree of the mamber of once patients.

At the request of the Calendar government, the factor of the perional Der doppent erran ed for epidendolosis add, outh a term of this national Der doppent erran ed for epidendolosis add, outh a term of this physicians from the Countries of the coloniant action action action action of the coloniant action of t

emainder of duiend. In the histogram, which appears on the lest page; asset of policeyelltis are thoen by data of passet of lines; or represe through January 15, as full analysa, of the epidemic will appear in a charquent issue of the PSU reports.


4. ENTEROVIRUS SURVEILLANCE

A total of 612 non-poliomyelitis enterovirus isolates from 1962 specimens have been reported from 34 States. Coxsackie B₃ was reported from 21 of these States, and ECHO₉ from 18 States. In California, Louisiana, Ohio and Texas, ECHO₉ accounted for the largest proportion of non-polio enteroviruses reported. In Connecticut, ECHO₄ accounted for the largest proportion of isolates made; in Hawaii - Coxsackie A₄ and A₅; in Michigan - Coxsackie A₉; and in New York - Coxsackie B₂.

5. FOREIGN REPORTS: British Guiana

Dr. B.B.G. Nehaul, Acting Chief Medical Officer for British Guiana has reported the occurrence of 361 cases of paralytic poliomyelitis through January 15, occurring in an epidemic which began during the last week in November. The age group involved has been predominantly the very young with 90 percent of the patients being below six years of age and almost 40 percent being below 2 years of abe. Type I poliovirus has been recovered from a large number of these patients.

At the request of the Guianian government, the Agency for International Development arranged for epidemiologic aid, with a team of four physicians from the Communicable Disease Center headed by Dr. Henry Gelfand to aid in the use of trivalent oral polic vaccine, supplied by the United States. This vaccine was administered beginning in Georgetown on December 22, with continued use during the following three weeks throughout the remainder of Guiana. In the histogram, which appears on the next page, cases of policyelitis are shown by date of onset of illness, as reported through January 15. A full analysis of the epidemic will appear in a subsequent issue of the PSU reports.



Non-Polio Enterovirus Isolations from 1962 Specimens

				ECHO				Coxs	ackie			
State		4	9	14	22	Other*	B-2	<u>B-3</u>	B-5	Other*	Total	Reported By
Arizona						1					1	T. Chin
Alabama						6				6	12	T. Hosty
California		6	24	2	6	9	4	4	1	16	72	E. Lennette and P. Wehrle
Connecticut		34	24	-	v	-	4 3	7	1 .7	10	54	E. Borman and G. Hsiung
Florida		2						1		2	5	J. Bond
Georgia						2		2	1	4	9	
Hawaii			1			2 2		1 7	7	36	46	J. Enright
Idaho			ĩ			-			1		1	D. Brack
			2			1	3	4	10.	4	14	H. Shaughnessy
lowa			-			6		-		2	8	T. Chin
Kansas	.5		1			11	4	10		9	35	C. Hunter
Kentucky			4	1			1	1	1	1	9	C. Todd
Louisiana			18	3		2		8		5	36	G. Hauser
Maryland	œ.		1	1		3	3	7		8	23	J. Joseph
Massachusetts						1		3		3	7	R. MacReady and J. Daniels
Michigan		3		6			1	5	7	10	32	G. Agate
Minnesota			5			11	3	18	-1	4	42	H. Bauer
Mississippi			2								2	H. Ricks
Missouri	18					11	1	4		3	19	I. Adams
Nebraska		*								1	- 1	T. Chin
Nevada			2						1		2	
New Hampshire	:								2		2	R. Miliner
New Jersey						1			3	1	5	M. Goldfield and W. Dougherty
New York	3.3		1	1	1	2	11	4	1	9	30	I. Albrecht
North Carolin	a		_	_	_	-			-	í	1	L. Madry
												-

Non-Polio Enterovirus Isolations from 1962 Specimens (Continued)

C+++	,	ЕСНО		\$ V.	Coxsackie	36°) 1 전 1 명 중 3 + 1 휴 7 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -
State 2 3	4	9 14	22 Other*	B-2	B-3 B-5	Other*	Total Reported By
Ohio		14	2 2 4	30	B & <	2	
Pennsylvania		2 1		10 E	11 4	7	43 C. Croft
Rhode Island	12	4 3 3 1 1 S	3 - 4 b 5 5	2	9 5 3		23 K. Hummeler and I. Gratch
Tennessee	•	1 4 6 6 7	음으로 번호		51		2 G. Hsiung
Texas	2	01.00	L 12 8 1 1 1		7	2	10 C. Tucker
Texas 4	0	21	2	2	1	ੁ 4	J. Irons and A. Behbehani
Utah			4 6 5 6	3 8	8 1. 8	3.	
Virginia		50 E	本音音 \$ \$		0.0,9	. 1	1 R. Fraser
Washington		1	3 8	- 10 - 11	2	2	7 E. Foxhall
Wisconsin		4	13 8 8 2 3 E		2	1 1	7 K. Berquist and W. Giedt
"15COH5III			3	1,	3	2	15 A. Evans
			* 0 0 0 0 0 0				
- 전 12, 10, 40		- 4 G 6	그 일 때 무 점				1 : x -

612

^{*} Specific types include 15 Coxsackie A-4 and 12 A-5 in Hawaii, and 6 A-9 in Michigan. A variety of single isolates of other ECHO and Coxsackie types have been reported.

STATEMENT OF COMMITTEE ON CONTROL OF INFECTIOUS DISEASES ON ORAL POLIOVACCINE

Released January 19, 1963

After careful re-evaluation of all previously reported cases and all newly reported cases of polio-like illness following administration of oral polio vaccine (OPV) in non-epidemic areas, Surgeon General Luther L. Terry and his Advisory Committee on Oral Poliomyelitis Vaccine recommended on December 18, 1962 that communities planning immunization campaigns against poliomyelitis "are urged to move ahead, using all three virus types of Sabin Oral Vaccine with particular emphasis on children and young adults." The Academy of Pediatrics Committee on Control of Infectious Diseases supports the recommendation and advises resumption of Type III OPV programs.

Excerpts from the Surgeon General's report which form the basis for the present Academy recommendation are as follows:

Some of the cases previously accepted as "compatible" have been dropped from this category because of additional clinical and laboratory evidence that has become available, and some new cases have been added.

The Committee does not consider that any individual case can be proved to be caused by the vaccine and no laboratory test has thus far provided a definite answer.

Considering the large amount of vaccine administered—it is concluded that contact spread to household contacts has posed no significant hazard.

As stated by the Surgeon General's committee, the maximum potential risk for Types I and III is of the order of one per million or less overall, but higher for those over 30 years of age. For Type II there is no indication of risk.

The Committee emphasized the importance of immunizing children in whom the danger of naturally occurring disease is greatest, and for adults who are at higher risk of naturally occurring disease; for example, parents of young children, pregnant women, persons in epidemic situations, and those planning foreign travel.

In summary: Two polio vaccines are now available and are recommended by the Public Health Service and endorsed by the Committee on Control of Infectious Diseases. Choice of which vaccine to use must rest with each community and each physician, and will be governed by local conditions, facilities, leadership, etc. Procedures for using Salk vaccine are well known. Procedures for using OPV should be guided by the recommendations of the Surgeon General and by the experience of numerous successfully

6.

completed community programs. The OPV is particularly indicated for Immunization of children who are the carriers of poliovirus in a community, for parents of young children because they are exposed to greater risk of natural infection, and in epidemic situations where it is more effective than killed vaccine.

Committee Members

Lewis L. Coriell, M.D., Chairman David Karzon, M.D. George A. McNaughton, M.D. Margaret H.D. Smith, M.D. Henry B. Strenge, M.D. Franklin H. Top, M.D. Dwain N. Walcher, M.D. Robert Ward, M.D. Thomas H. Weller, M.D. Sarah H. Wood Sell, M.D. Guillermo Tovar, M.D.

Figure 1a CURRENT U.S. POLIO INCIDENCE COMPARED WITH YEARS 1957, 1959 and 1961 April-December, by week




Figure 1

CURRENT U.S. POLIO INCIDENCE
COMPARED WITH YEARS 1958, 1960, and 1962




Table 1
TREND OF 1963 POLIOMYELITIS INCIDENCE

and Region UNITED STATES Paralytic Nonparalytic	1963 6	12/8	For 12/15	72/22				Week	Week	Totals	in
UNITED STATES Paralytic Nonparalytic		12/8	12/15	72/22							
Paralytic Nonparalytic	6			14/44	12/29	1/5	1/12	Total	1962	1961	1960
Paralytic Nonparalytic	6										
Nonparalytic		5	5	18	8	1	5	42	56	123	282
	1	2	ĭ	2	2	_	ì	8	22	29	64
Unspecified	_		_	_	_	-	_	_	23	53	29
Total	7	7	6	20	10	1	6	50	101	205	375
NEW ENGLAND											
Paralytic	_	_	_		-		_	_	4	4	16
Total	_	-	_	_	-	-		_	6	6	17
Maine	ř	_	_	-			_	_	2	4	6
New Hampshire	_	_	_	~	_	_	_	_	ĩ	7	_
Vermont	-	_	_	_	-	_	_	_	ī	7	1
Massachusetts	- <u>-</u> \	_	_	_	_	-	_	_	2	1 <	7
Rhode Island	-	-	_	_	_	_	_	_		ī	
Connecticut	/ = -	_	_	-	_	_	_	_	· -	-	3
MIDDLE ATLANTIC											
Paralytic	_	_	1	5	2	_		8	12	27	38
Total		_	ī	5	2		_	8	24	44	54
New York		_	ī	_		_	_	ĭ	22	26	38
New Jersey		_	-	_		_	_		- I	2	4
Pennsylvania	-	-	-	5	2	-	-	7	2	16	12
EAST NORTH CENTRAL	L										
Paralytic	2	3	1	4		-	2	10	2	25	40
Total	3	3	1	5			3	12	6	34	68
Ohio	ĭ	_	_	_	-	_	ì	1	1	5	22
Indiana	1 Sec.	2		1		_	_	3	3	13	4
Illinois	2	_		4	_		2	6	ì	9	32
Michigan	_ <u></u>	1	_	- 10	_	_		ì	ī	2	6
Wisconsin	4 ⁻	_	1	- ,	-	-	-	ī	_	5	4
WEST NORTH CENTRAL	L										
Paralytic	7-4		_	3	_	_	_	3	3	5	15
Total		_		3	-	_	_	3	10	7	27
Minnesota	_	_	_	_	_		_	-	2	1	3
Iowa		_	-	-	_		_	_	2	1	_
Missouri	1	_	_	2	_	_	_	2	4	1	9
North Dakota			_	_	_	_	-	-	ì	2	_
South Dakota		1		ī	-	***	_	1	ī	ī	_
Nebraska	_		14.	_		_	_	_	_	ī	1
Kansas	-		_	_	_	_	_	_	_	_	8

Table 1 (Continued)

State and	Cumula- tive	•	Cases For	Report Week	ted to Ending	CDC		Six Week	Compa Weeks	rable Total	
Region	1963	12/8	12/15			1/5	1/12	Total	1962	1961	1960
COLUMN A MY ANIMY O											
SOUTH ATLANTIC					7			7	~	26	48
Paralytic Total	-	-	-	-	1 2	-	-	1 2	7	26	
Delaware	-	-	_	_		-	-		9	32	59
Maryland	-	-	-	_	-	-	-	-	4	$\begin{array}{c} 1 \\ 12 \end{array}$	2
D.C.	_	-		_	-	-	_	-			í
Virginia	-	_	_	_	-	_		_	-	7	4
West Virginia	_	_	_	-	_		_	_	2	í	6
North Carolina	_	_	_	_		_	_		_	3	22
South Carolina	_	_	_	_	-	_	_	-		2	1
Georgia	_	_	-	-	2	-	-	2	-	2	17
Florida	_	_	-	_	_	-	-		3	4	6
riorida	-	-	-	-	-	-		-	3	4	O
EAST SOUTH CENTRA	AL							,			
Paralytic	-	_	-	2	1	_	-	3	5	5	24
Total	-	-	-	2	1	_	_	3	5	31	28
Kentucky	_	_	-	1	-	-	_	1	_	25	9
Tennessee	_	-	-	_	_	-	-	_	-	3	7
Alabama	-	_	-	-	1	-	_	1	2	1	2
Mississippi	-	-	-	1	_	_	_	1	3	2	10
WEST SOUTH CENTRA	Λ Τ.										
			2	0	0		0	^	10	11	18
Paralytic	2	-	3 4	2 3	2	-	2 2	9 12	10 12	15	24
Total Arkansas	2	_			3	-			2		
Touris	_	-	2	_	_	-	2	~	2	6 5	5 7
Louisiana	2	_		2	3	-		9			6
Oklahoma Texas	_	-	1 1	ī	_	-	-	1 2	- 8	- 4	6
CAAS	-	_	1	1	_	-	_	2	0	*	U
MOUNTAIN											
Paralytic	-	-	-	-	-	-	_	-	6	4	9
Total	-	-	-	-	_	-	-	-	17	11	12
Montana	_	-	- ,	_	_	-	-	_	2	2	5
Idaho	-	_	_	-	_	-	-	-	1	2	-
Wyoming	_	-	-		-	-	-	-	1	1	-
Colorado	_	-	-	-	-	-	-		1	-	-
New Mexico	-	-	-	-	-	-	-	-	10	-	1
Arizona	-	-	-	-	-	-	-	-	1	1	6
Utah	-	-	-	-		-	-	-	-	5	-
Nevada	-	-	-	-	_	-	-	-	1	_	-
PACIFIC	2	2		2	2	1	1	8	7	16	74
Paralytic	2	2	_	2	2	ì	ì	10	12	25	86
Total	2	4	-			T		10	12	10	17
Washington	-	-	-	-	_	-	_	_			
Oregon		-	-	-	_	-	-	7.0	- 3.0	2	21
California	2	4	_	2	2	1	1	10	10	13	46
Alaska	-	-	-		-	-	-	-	-	-	1
Hawaii	-	-	-	-	-	-	-	-	1	-	1
TERRITORY											
Puerto Rico	-	_	_	1	_	_	_	1		2	7
OI CO KICO											

6	-alumuQ -vij				ted to Ending			Nist Week		rable Six
	1963	0101	A I'V	- A C F	ladini 12/29	- diese	-	24515 W		
And the second s	COGT	0 /24	Carlotte State	and the second	27 / 1977			1830)	1962	1901 1901
OTTHAITA I										
pingism					Ω					A
isi								2	1	26 48
9.18				79-				:2	ē	32 59
and										- L
W. C. & F. W.	-			1.40	8-2					1.2
sin								••		1
Virgan	,,,,,,,,						-	*	2	
Car oldina.				46			140			
Carolina										50 C
ia	***						11-89.	2		2 1
da	***						***	10		2 17 4 6
			44.5	4,5				4-2	3	0 4
SOUTH CENTURE										
ralytic			Acres .	1	ſ			3	***	5 24
ist				2	Ĭ.		-	8	13.7-8	
oks	940 J-	Poli	***	Ĩ						6 25
ssee		Specif		-				.5-		7.5.5.6
£m		(in)						1	2	2
- iqqiee	197.6		146	3				Ī		2 10
								1,		
SOUTH CENTRAL										
ralgtic V	25		- 8	5			S	ė	- 01	93 18
tal		***	4	£.			3	SI	2.1	15 24
80.8			979				144	W.A.	2	
ensl			6	. 5				6	2	7 - 3
oma		N. 70	T I	1	700			1	-	a –
	100		Ĺ	1			-		8	3 k
				wire.						
9 MIA										
ralycic	1.0	70.		ξ		***	4940	1		6 4
1 Hst	***			200	1960	8-10	100		1.7	of It
3 50							pen.		2	2 65
				***			Sec.	(6)	Ī	2 11 11 12
1 20	140		200	14		-		417	Ī	
obs	***		- 10	-4-			1994		- 1	
end.co	Supr.	1077	447	0.00		-		_		1
	to the second			To the same of the						a l
	-				100			Topic Control		
10 7 1						4.50	Marie			- 10
01										
oidAla	2		-	5	S	I	1			15 24
fas	2			2	2	j	1	0.0		25 86
neton	And .	-		1000			1946	-		71 01
- 1				,		4.0				2 21
rnia	2	20		2	2		I	ŌΙ	10	0 17 2 21 13 46
	144									Parameter and St.

SUPPLEMENT TO

Poliomyelitis Surveillance Report No. 274

January 22, 1963

INDEX TO 1962
POLIOMYELITIS SURVEILLANCE REPORTS
(NOS. 249 - 273)

OT TIMELIPHUE

Foliomyelitis Surveillance Report No. 274

January 22, 1963

INDEX TO 1962 FOLIOMYFLITIS SURVEILLANCE REPORTS (NOS. 249 - 273)

INDEX TO 1962 PSU REPORTS (NOS. 249-273)

nten	its		PSU	Repor	t Num	ber	<u>on</u>
cu	RRENT POLIOMYELITIS				SICI	, е	
Α.	Reports From States	Republica Valers by	(Variety				
•••	Alabama Double	Recail acidedicost					
	Arizona	white and Vaccinetics	252		1 ve 4		
	Arkansas			8.03 8			
	California				267,	268	2.72
	Illinois	STANLE COFFEE			3/11		212
	Vantualar		266	267			
0	Massachusetts	OSY to evon-Or what	261				
	Minnesota						
	Nebraska		268	271			
	New York State		249,	250			
	Ohio	Ancirsts of Paralyth	252	200	1001		
	Oklahoma						
	Pennsylvania			266			
	Toyas	Day to suell-06 aids!					
	Virginia	Total	266	200,	203	4.5	
	West Virginia	Prograf Lauft La					
	West Vilginia		daoqeA :				
В.	Foreign Reports				0000		
ь.	Argentina		253				
	Australia	L Vaucineas	272				
	Canada		251,				
	Chile	real and in third are	249.	250.	261		
	England		250	,			
	Israel		254				
	Japan			252			
366	Philippines	e Distribution					
268	Tacamology 2. V21		,				
OR.	AL POLIOVIRUS VACCIN	NATION PROGRAMS					
	Atlanta, Georgia		249,	253	$rd \in K$		
	Newberry, South Car	rolina	249,	253			
	Tampa-Hillsborough	County, Florida				av.	
	Mass Oral Polio Vac		271				
	180 181 181				Legi		
PO	LIOMYELITIS CASES RE	EPORTED TO PSU					
Α.	1961				1143	3.6	
	Cases by Age and Va		249-				
		e and Vaccination					
	History (1961)		254				
		ities and Age-Specific					
	Attack Rates (19		255				
	Final Report-Type of						
	Involvement (196		256				
		on by Virus Isolations					
	(1961)		260				

INDEX TO 1962 PSU REPORTS (CONTINUED)

	PSU Report Must	PSU Report Number
В.	1962 Eva Company Reported Cases by	1. CURRENT POLICHYBLIT
	County (Map)	253255, 257 ₄
	Paralytic Cases by Vaccination Status	259, 261, 262
	Paralytic Cases by Age and Vaccination	2012 CHAN 60 64 751 7511/2
	Status	269 273
000		268273 ARNA
	256, 263, 267,	California
. <u>RO</u>	UTINE POLIOMYELITIS SURVEILLANCE	aiontLII
Α.	Cases with Onset Within 30-Days of Vaccin	ation (Inactivated)
		249251
	1961 Cases Reported	
	1961 Cases Reported-Final Report	253 Salasadon
	1962 Cases Reported	254273
	1961-1962 Summary Analysis of Paralytic	ohiio
	Cases ETS	270 emoris (510
	250, 266	Remarkania Pennetina
В.	Cases with Onset Within 30-Days of Vaccin	ation (Oral)
	1961 Cases Reported	249251
	1961 Cases Reported-Final Report	253 regret rest
	1962 Cases Reported	254273
	1902 Cases Reported	
c.	Polio Within 60-Days Among Household	B. Foreign Reports
•	Contact of Oral Vaccinees	271273
	OSSO PSO	
n	TORO A DORO DITTO A CONTRACTOR AND A CON	Canada
D.	Inactivated Vaccine Distribution	Chile
	1961 Inactivated Vaccine Distribution	249251
	1962 Inactivated Vaccine Distribution	252, 253, 255, 257
170	Our Delia Wooding Distribution	PCU Sample 266
Ε.	Oral Polio Vaccine Distribution	PSU Supplement 266
		PSU Supplement 268
	CONNETTION PROCESSES	PSU Supplement 273
	Nebraska	Atlanta, 600 gil
***	NEW YORK MYALL BEIDAD MEIN MA DOLL	Newberry, South
\cdot vi		Tampa-Hillsberou
	Vaccination Programs 271	Mass Oral Ped
	1961 Poliovirus Isolates by State	250, 251, 254
		259273
ΔQ	1962 Poliovirus Isolates by State 1904 AR	250 273
. <u>AS</u>	1962 Poliovirus Isolates by State 190434 a	2. 1961 .A
. AS	1962 Poliovirus Isolates by State 190434 a EPTIC MENINGITIS 232042 200832 noidenioneV b	2. FOLLOWER 273 AVEC AND ARREST ARRES
. <u>AS</u>	1962 Poliovirus Isolates by State 190434 a	249, 251, 252, 254, 257
. AS	1962 Poliovirus Isolates by State 190434 & EPTIC MENINGITIS Nation-wide Enterovirus Surveillance 1944	249, 251, 252, 254, 257 260, 267, 270272
. <u>AS</u>	1962 Poliovirus Isolates by State 190434 & EPTIC MENINGITIS Nation-wide Enterovirus Surveillance 1944	249, 251, 252, 254, 257 260, 267, 270272
. <u>AS</u>	1962 Poliovirus Isolates by State 190439 & EPTIC MENINGITIS Nation-wide Enterovirus Surveillance and Connecticut offices and beautifus and connecticut offices an	249, 251, 252, 254, 257 260, 267, 270272 252, 269
. <u>AS</u>	1962 Poliovirus Isolates by State 190439 & EPTIC MENINGITIS Nation-wide Enterovirus Surveillance 1904 (1904) Connecticut 1904 (1904)	259273 249, 251, 252, 254, 257 260, 267, 270272 252, 269 271
. <u>AS</u>	1962 Poliovirus Isolates by State 190439 & EPTIC MENINGITIS Nation-wide Enterovirus Surveillance SA Connecticut Minnesota 122 (1901) Texas	249, 251, 252, 254, 257 260, 267, 270-272 252, 269 271 249, 269
. AS	Poliovirus Isolates by State 190439 & EPTIC MENINGITIS Nation-wide Enterovirus Surveillance and Minnesota (1901) Texas (1902)	249, 251, 252, 254, 257 260, 267, 270272 252, 269 271 249, 269
. <u>AS</u>	1962 Poliovirus Isolates by State 190439 & EPTIC MENINGITIS Nation-wide Enterovirus Surveillance SA Connecticut Minnesota 122 (1901) Texas	249, 251, 252, 254, 257 260, 267, 270272 252, 269 271 249, 269

INDEX TO 1962 PSU REPORTS (CONTINUED)

Contents	PSU R	eport Nu	mber
7. MISCELLANEOUS		LEAT .	1.0
"Recommendations of the Public Health Servi for the Use of Poliomyelitis Vaccines for the 1962 Poliomyelitis Season" Intratypic Sero-differentiation of Poliovir Special Report Revisions in Poliomyelitis Surveillance Cas Record News Release-October 2, 1962	253 ruses- 256	71 47 80	
8. SUPPLEMENTS			3.1
Index to 1961 PSU Reports Poliomyelitis Vaccination Survey-Syracuse, New York-November 1961 Poliomyelitis Vaccination Survey-San Juan, Puerto Rico-May 1961 Special Report-Cases of Poliomyelitis Occur Within 30 Days Following Oral Vaccine Administration-August 1962 A. Surgeon General's News Release B. Background Material C. Case Summaries and Oral Vaccine Usage Surgeon General's Technical Report to the M Profession Concurring Oral Vaccine Usage "The Association of Cases of Poliomyelitis the Use of Type III Oral Poliomyelitis V Case Summaries of Poliomyelitis Associated Administration of Oral Vaccine-Cases Comby Surgeon General's Committee through O	254 255 rring 266 26e 26e 26e 26e 26e 26e 26e 26e 26e		
News Release following Surgeon General's OF Advisory Committee Meeting-December 17, Report of the Special Advisory Committee on Poliomyelitis Vaccine to the Surgeon Gen Cases Considered by the Special Advisory Co	1962 273 n Oral neral, PHS 273		
9. <u>FIGURES</u>			
Current U.S. Polio Incidence Quarterly Shipment of Poliomyelitis Vaccine (Inactivated) 1955-1961 Current Polio Incidence in Japan Poliomyelitis Cases by Age, Tokyo, Japan: Paralytic Poliomyelitis Cases by Week of On	251 251 1960-61 252 aset-	73	
Texas, 1962	269		

INDEX TO 1962 PSU REPORTS (CONTINUED)

Cont	ents:e8 USS	PS	U Repor	t Num	iber
10.	TABLES	210	CELLANE	2114	7.
	Trends of 1961 Poliomyelitis Incidence	moiseb ₂₄	ono se Hi		
	Trends of 1962 Poliomyelitis Incidence	980 925	0273		
	Vaccine Distribution "posses similaymoi	169 2024	9253	255	2.57
	Quarterly Shipments of Poliomyelitis Vaccine;	ole Sere	Intraty	200,	201
	1955-1961	rogeA [25	Spe		
	Mass Oral Polio Vaccination Programs from Month				
	of Vaccine Licensure through October 10, 196		Reco		
	cober 2, 1362 - 269 -				
11.	MAPS				
	* Marianta M	3	THEMENT	172	. 8
	Australia-Polio Case Attack Rate	27:	CONTRACTOR NAME OF THE PARTY OF		
	Nebraska-Counties in which Type III OPV was				
	Administered Same and Administered	7 27	Politians		
	San Juan, Puerto Rico-Identification of Survey				
	Districts				
	San Juan, Puerto Rico-Socio-economic Area				
	Classification of the Survey Districts				
	Syracuse, New York ogioos land native for an				
	Reported Paralytic Poliomyelitis in Texas by				
				000	
	County, 1962 esseled even a largest Vaccine Usage in U.S.			269	
	marios and Oral Vaccine Usage				
	's Technical Report to the Medical				
	oncurring wral Accoine Usages				
	tion of Cares of Poliomyelitis with				
	ype III Oral Poliomyelitis Vaccines 268				
	of Poliomyelitiz associated with the				
	on of Oral Vaccine-Cases Considered				
	eneral's Committee through October 2,				
	270		1961		
	Llowing Surgeon General's OPV				
	mittee Meeting-December 17, 1962 273				
	pecial Advisory Committee on Oral				
	s Vaccine to the Surgeon General, 1915 273	itilsyno.	t Log		
	d by the Special Advisory Committee 323	onsidare	Cases C		
			EHAD.		
	Lio Incidence 249-273	. U.S. Po	Current		
	ept of Policeyelitis Vaccine	mqiris vi	Quar cer		
) 1955-1961				
	neidence in Japan 251				
	again by Aga Polyna Jayan 1666 61 282	n what			